

HOMWORK ASSIGNMENT #2, Math 253

- Find the equation of a sphere if one of its diameters has end points $(1, 0, 5)$ and $(5, -4, 7)$.
- Find vector, parametric, and symmetric equations of the following lines.
 - the line passing through the points $(3, 1, \frac{1}{2})$ and $(4, -3, 3)$
 - the line passing through the origin and perpendicular to the plane $2x - 4y = 9$
 - the line lying on the planes $x + y - z = 2$ and $3x - 4y + 5z = 6$
- Find the equation of the following planes.
 - the plane passing through the points $(-1, 1, -1)$, $(1, -1, 2)$, and $(4, 0, 3)$
 - the plane passing through the point $(0, 1, 2)$ and containing the line $x = y = z$
 - the plane containing the lines

$$L_1 : x = 1 + t, \quad y = 2 - t, \quad z = 4t$$

$$L_2 : x = 2 - s, \quad y = 1 + 2s, \quad z = 4 + s$$

- Find the intersection of the line $x = t, y = 2t, z = 3t$, and the plane $x + y + z = 1$.
- Find the distance between the point $(2, 8, 5)$ and the plane $x - 2y - 2z = 1$.
- Show that the lines

$$L_1 : \frac{x - 4}{2} = \frac{y + 5}{4} = \frac{z - 1}{-3}$$

$$L_2 : \frac{x - 2}{1} = \frac{y + 1}{3} = \frac{z}{2}$$

are skew. Find the distance between the two lines.

- Identify and sketch the following surfaces.
 - $4x^2 + 9y^2 + 36z^2 = 36$
 - $4z^2 - x^2 - y^2 = 1$
 - $y^2 = x^2 + z^2$
 - $x^2 + 4z^2 - y = 0$
 - $y^2 + 9z^2 = 9$
 - $y = z^2 - x^2$
- Find the polar equation for the curve represented by the following Cartesian equation.
 - $x = 4$

(b) $x^2 + y^2 = -2x$

(c) $x^2 - y^2 = 1$

9. Sketch the curve of the following polar equations.

(a) $r = 5$

(b) $\theta = \frac{3\pi}{4}$

(c) $r = 2 \sin \theta$

(d) $r = 3(1 - \cos \theta)$

10. (a) Change $(3, \frac{\pi}{3}, 1)$ from cylindrical to rectangular coordinates

(b) Change $(\sqrt{3}, 1, 4)$ from rectangular to cylindrical coordinates

(c) Change $(\sqrt{3}, 1, 2\sqrt{3})$ from rectangular to spherical coordinates

(d) Change $(4, \frac{\pi}{4}, \frac{\pi}{3})$ from spherical to cylindrical coordinates

SOLUTIONS TO HOMEWORK ASSIGNMENT #2, Math 253

1. Find the equation of a sphere if one of its diameters has end points $(1, 0, 5)$ and $(5, -4, 7)$.

Solution:

The length of the diameter is $\sqrt{(5-1)^2 + (-4-0)^2 + (7-5)^2} = \sqrt{36} = 6$, so the radius is 3. The centre is at the midpoint $(\frac{1+5}{2}, \frac{0-4}{2}, \frac{5+7}{2}) = (3, -2, 6)$. Hence, the sphere is given as $(x-3)^2 + (y+2)^2 + (z-6)^2 = 9$.

2. Find vector, parametric, and symmetric equations of the following lines.

- (a) the line passing through the points $(3, 1, \frac{1}{2})$ and $(4, -3, 3)$

Solution:

The vector between two points is $\vec{v} = \langle 4-3, -3-1, 3-\frac{1}{2} \rangle = \langle 1, -4, \frac{5}{2} \rangle$. Hence the equation of the line is

Vector form: $\vec{r} = \vec{r}_0 + t\vec{v} = \langle 4, -3, 3 \rangle + t\langle 1, -4, \frac{5}{2} \rangle = \langle 4+t, -3-4t, 3+\frac{5}{2}t \rangle$

Parametric form: $x = 4+t, \quad y = -3-4t, \quad z = 3+\frac{5}{2}t$

Symmetric form: Solving the parametric form for t gives $x-4 = \frac{y+3}{-4} = \frac{z-3}{5/2}$

- (b) the line passing through the origin and perpendicular to the plane $2x - 4y = 9$

Solution:

Perpendicular to the plane \Rightarrow parallel to the normal vector $\vec{n} = \langle 2, -4, 0 \rangle$. Hence

Vector form: $\vec{r} = \langle 0, 0, 0 \rangle + t\langle 2, -4, 0 \rangle = \langle 2t, -4t, 0 \rangle$

Parametric form: $x = 2t, \quad y = -4t, \quad z = 0$

Symmetric form $\frac{x}{2} = \frac{y}{-4}, \quad z = 0$

- (c) the line lying on the planes $x + y - z = 2$ and $3x - 4y + 5z = 6$

Solution:

We can find the intersection (the line) of the two planes by solving z in terms of x , and in terms of y .

$$(1) \quad x + y - z = 2$$

$$(2) \quad 3x - 4y + 5z = 6$$

Solve z in terms of y : $3 \times (1) - (2) \Rightarrow 7y - 8z = 0 \Rightarrow z = \frac{7}{8}y$

Solve z in terms of x : $4 \times (1) + (2) \Rightarrow 7x + z = 14 \Rightarrow z = 14 - 7x$

Hence, symmetric form: $14 - 7x = \frac{7}{8}y = z$

Set the symmetric form $= t$, we have parametric form: $x = \frac{14-t}{7}, \quad y = \frac{8}{7}t, \quad z = t$

Vector form: $\vec{r} = \langle \frac{14-t}{7}, \frac{8}{7}t, t \rangle$

3. Find the equation of the following planes.

- (a) the plane passing through the points $(-1, 1, -1)$, $(1, -1, 2)$, and $(4, 0, 3)$

Solution:

Name the points $P(-1, 1, -1)$, $Q(1, -1, 2)$, and $R(4, 0, 3)$. Set up two vectors:

$$\vec{v}_1 = \overrightarrow{PQ} = \langle 1 + 1, -1 - 1, 2 + 1 \rangle = \langle 2, -2, 3 \rangle \quad (1)$$

$$\vec{v}_2 = \overrightarrow{PR} = \langle 5, -1, 4 \rangle \quad (2)$$

Choose the normal vector $\vec{n} = \vec{v}_1 \times \vec{v}_2 = \langle -5, 7, 8 \rangle$. Hence the equation of the plane is $\boxed{-5(x + 1) + 7(y - 1) + 4(z + 1) = 0}$ using point P .

- (b) the plane passing through the point $(0, 1, 2)$ and containing the line $x = y = z$

Solution:

Name $Q(0, 1, 2)$. The line can be represented as $\vec{r} = \langle t, t, t \rangle$, which crosses the point $P(0, 0, 0)$ and is parallel to $\vec{v} = \langle 1, 1, 1 \rangle$. Set $\vec{b} = \overrightarrow{PQ} = \langle 0, 1, 2 \rangle$. Choose $\vec{n} = \vec{v} \times \vec{b} = \langle 1, -2, 1 \rangle$ and hence the equation of the plane is $\boxed{x - 2y + z = 0}$ using point P .

- (c) the plane containing the lines

$$L_1 : x = 1 + t, \quad y = 2 - t, \quad z = 4t$$

$$L_2 : x = 2 - s, \quad y = 1 + 2s, \quad z = 4 + s$$

Solution:

From L_1 and L_2 , $\vec{v}_1 = \langle 1, -1, 4 \rangle$ and $\vec{v}_2 = \langle -1, 2, 1 \rangle$. Choose $\vec{n} = \vec{v}_1 \times \vec{v}_2 = \langle -9, -5, 1 \rangle$. Since L_1 crosses the point $(1, 2, 0)$, the equation of the plane is $\boxed{-9(x - 1) - 5(y - 2) + z = 0}$.

4. Find the intersection of the line $x = t$, $y = 2t$, $z = 3t$, and the plane $x + y + z = 1$.

Solution:

Substitute the line into the plane: $t + 2t + 3t = 1 \Rightarrow t = \frac{1}{6}$.

Put t back to the line: $x = \frac{1}{6}$, $y = \frac{1}{3}$, $z = \frac{1}{2}$.

Hence the intersection point is $\boxed{(\frac{1}{6}, \frac{1}{3}, \frac{1}{2})}$.

5. Find the distance between the point $(2, 8, 5)$ and the plane $x - 2y - 2z = 1$.

Solution:

Name $Q(2, 8, 5)$. Choose any point on the plane, say a convenient one $(x, 0, 0)$. So $x - 2(0) - 2(0) = 1 \Rightarrow x = 1 \Rightarrow P(1, 0, 0)$. Then $\vec{b} = \overrightarrow{PQ} = \langle 1, 8, 5 \rangle$. The normal vector of the plane is $\vec{n} = \langle 1, -2, -2 \rangle$. The distance between the plane and the point is given as

$$\text{distance} = \left| \text{proj}_{\vec{n}} \vec{b} \right| = \frac{|\vec{n} \cdot \vec{b}|}{|\vec{n}|} = \frac{|-25|}{|3|} = \boxed{\frac{25}{3}}$$

6. Show that the lines

$$L_1 : \frac{x-4}{2} = \frac{y+5}{4} = \frac{z-1}{-3}$$

$$L_2 : \frac{x-2}{1} = \frac{y+1}{3} = \frac{z}{2}$$

are skew.

Solution:

Write the equation in parametric form.

$$L_1 : x = 2t + 4, \quad y = 4t - 5, \quad z = -3t + 1$$

$$L_2 : x = s + 2, \quad y = 3s - 1, \quad z = 2s$$

The lines are not parallel since the vectors $\vec{v}_1 = \langle 2, 4, -3 \rangle$ and $\vec{v}_2 = \langle 1, 3, 2 \rangle$ are not parallel. Next we try to find intersection point by equating x , y , and z .

$$(1) \quad 2t + 4 = s + 2$$

$$(2) \quad 4t - 5 = 3s - 1$$

$$(3) \quad -3t + 1 = 2s$$

(1) gives $s = 2t + 2$. Substituting into (2) gives $4t - 5 = 3(2t + 2) - 1 \Rightarrow t = -5$. Then $s = -8$. However, this contradicts with (3). So there is no solution for s and t . Since the two lines are neither parallel nor intersecting, they are skew lines.

7. Identify and sketch the following surfaces.

(a) $4x^2 + 9y^2 + 36z^2 = 36$

Solution:

xy -plane: $4x^2 + 9y^2 = 36$ ellipse

xz -plane: $4x^2 + 36z^2 = 36$ ellipse

yz -plane: $9y^2 + 36z^2 = 36$ ellipse

\Rightarrow ellipsoid

(b) $4z^2 - x^2 - y^2 = 1$

Solution:

xy -plane: $-x^2 - y^2 = 1$ nothing, try $z = \text{constants}$

$z = c$: $-x^2 - y^2 = 1 - 4c^2 \Rightarrow x^2 + y^2 = 4c^2 - 1$ circles when $4c^2 - 1 > 0$

xz -plane: $4z^2 - x^2 = 1$ hyperbola opening in z -direction

yz -plane: $4z^2 - y^2 = 1$ hyperbola opening in z -direction

\Rightarrow hyperboloid of two sheets

(c) $y^2 = x^2 + z^2$

Solution:

xy -plane: $y^2 = x^2$ cross

xz -plane: $0 = x^2 + z^2$ point at origin, try $y = \text{constants}$

$y = c$: $c^2 = x^2 + z^2$ circles

yz -plane: $y^2 = z^2$ cross

\Rightarrow cone

(d) $x^2 + 4z^2 - y = 0$

Solution:

xy -plane: $x^2 - y = 0 \Rightarrow y = x^2$ parabola opening in $+y$ -direction

xz -plane: $x^2 + 4z^2 = 0$ point at origin, try $y = \text{constants}$

$y = c$: $x^2 + 4z^2 - c = 0 \Rightarrow x^2 + 4z^2 = c$ ellipses when $c > 0$

yz -plane: $4z^2 - y = 0 \Rightarrow y = 4z^2$ parabola opening in $+y$ -direction

\Rightarrow elliptic paraboloid

(e) $y^2 + 9z^2 = 9$

Solution:

x missing: cylinder along x -direction

yz -plane: $y^2 + 9z^2 = 9$ ellipse

\Rightarrow elliptic cylinder

(f) $y = z^2 - x^2$

Solution:

xy -plane: $y = z^2$ parabola opening in $+y$ -direction

xz -plane: $0 = z^2 - x^2 \Rightarrow z^2 = x^2$ cross, try $y = \text{constants}$

$y = c$: $c = z^2 - x^2$ hyperbola opening in z -direction when $c > 0$, in x -direction when $c < 0$

yz -plane: $y = -x^2$ parabola opening in $-y$ -direction

\Rightarrow hyperbolic paraboloid

8. Find the polar equation for the curve represented by the following Cartesian equation.

(a) $x = 4$

Solution:

$x = 4 \Rightarrow r \cos \theta = 4 \Rightarrow$ $r = 4 \sec \theta$

(b) $x^2 + y^2 = -2x$

Solution:

$x^2 + y^2 = -2x \Rightarrow r^2 = -2r \cos \theta \Rightarrow$ $r = -2 \cos \theta$




Figure 1: Q7(a)




Figure 2: Q7(b)




Figure 3: Q7(c)




Figure 4: Q7(d)




Figure 5: Q7(e)




Figure 6: Q7(f)

(c) $x^2 - y^2 = 1$

Solution:

$$x^2 - y^2 = 1 \Rightarrow r^2 \cos^2 \theta - r^2 \sin^2 \theta = 1 \Rightarrow r^2 (\cos^2 \theta - \sin^2 \theta) = 1 \Rightarrow r^2 \cos 2\theta = 1$$

$$\Rightarrow r^2 = \sec 2\theta \Rightarrow \boxed{r = \pm \sqrt{\sec 2\theta}}$$

9. Sketch the curve of the following polar equations.

(a) $r = 5$

(b) $\theta = \frac{3\pi}{4}$

(c) $r = 2 \sin \theta$

(d) $r = 3(1 - \cos \theta)$

10. (a) Change $(3, \frac{\pi}{3}, 1)$ from cylindrical to rectangular coordinates

Solution:

$$x = r \cos \theta = 3 \cos \frac{\pi}{3} = \frac{3}{2}, \quad y = r \sin \theta = 3 \sin \frac{\pi}{3} = \frac{3\sqrt{3}}{2}, \quad z = 1. \quad \text{Hence } (x, y, z) =$$

$$\boxed{\left(\frac{3}{2}, \frac{3\sqrt{3}}{2}, 1\right)}$$

(b) Change $(\sqrt{3}, 1, 4)$ from rectangular to cylindrical coordinates

Solution:

$$r = \sqrt{x^2 + y^2} = \sqrt{3 + 1} = 2, \quad \tan \theta = \frac{y}{x} = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6} \text{ in first quadrant, } z = 4.$$

$$\text{Hence } (r, \theta, z) = \boxed{\left(2, \frac{\pi}{6}, 4\right)}$$




Figure 7: Q9(a)




Figure 8: Q9(b)




Figure 9: Q9(c)




Figure 10: Q9(d)

- (c) Change $(\sqrt{3}, 1, 2\sqrt{3})$ from rectangular to spherical coordinates

Solution:

$\rho = \sqrt{x^2 + y^2 + z^2} = \sqrt{3 + 1 + 12} = 4$, $\tan \theta = \frac{y}{x} = \frac{1}{\sqrt{3}} \Rightarrow \theta = \frac{\pi}{6}$ in first quadrant, $\phi = \cos^{-1} \frac{z}{\rho} = \cos^{-1} \frac{2\sqrt{3}}{4} = \cos^{-1} \frac{\sqrt{3}}{2} = \frac{\pi}{6}$. Hence $(\rho, \theta, \phi) = \boxed{(4, \frac{\pi}{6}, \frac{\pi}{6})}$

- (d) Change $(4, \frac{\pi}{4}, \frac{\pi}{3})$ from spherical to cylindrical coordinates

Solution:

$r = \rho \sin \phi = 4 \sin \frac{\pi}{3} = 2\sqrt{3}$, $\theta = \frac{\pi}{4}$, $z = \rho \cos \phi = 4 \cos \frac{\pi}{3} = 2$. Hence $(r, \theta, z) = \boxed{(2\sqrt{3}, \frac{\pi}{4}, 2)}$