

HOMWORK ASSIGNMENT #7

- The iterated integral $I = \int_{x=0}^{x=1} \left(\int_{y=0}^{y=\sqrt{x}} \sin \left(\frac{\pi(y^3 - 3y)}{2} \right) dy \right) dx$ is equal to the double integral $\iint_R \sin \left(\frac{\pi(y^3 - 3y)}{2} \right) dA$ for a region R in the x, y plane.
 - Sketch R .
 - Write the integral with the order of integration reversed.
 - Compute I .
- Let D be the region bounded by $y = x$ and $y = 6 - x^2$.
 - Sketch D .
 - Find $\iint_D x^2 dA$.
- Let D be the region, described in polar coordinates by, $0 \leq \theta \leq \pi, 0 \leq r \leq 1 + \cos \theta$.
 - Sketch D .
 - Compute the area of D .
 - Find the average value of distances of points in D from the origin.
- Determine the following integrals:
 - $\iint_D (|x| + |y|) dA$, where D is the region $x^2 + y^2 \leq a^2$ and a is a positive constant.
 - $\iint_T \sqrt{a^2 - x^2} dA$, where T is the triangle with vertices $(0, 0), (a, 0), (a, a)$.
 - $\iint_D \frac{1}{x^2 + y^2} dA$, where D is the region in the first quadrant bounded by
$$y = 0, y = x, x^2 + y^2 = 1/4, x^2 + y^2 = 1.$$
 - $\iint_R (\sin xy + x^2 - y^2 + 3) dx dy$, where R is the region inside the circle $x^2 + y^2 = a^2$ and outside the circle $x^2 + y^2 = b^2$, and a, b are constants satisfying $0 < b < a$.
- Find the volume above the x, y plane, below the surface $z = e^{-(x^2+y^2)}$ and inside the cylinder $x^2 + y^2 = 4$.
- Find the volume above the x, y plane and below the surface $z = e^{-(x^2+y^2)}$.
- Compute the double integral $\iint_D (x + y) dA$, where D is the domain that lies to the right of the y -axis and between the circles $x^2 + y^2 = 1, x^2 + y^2 = 4$.
- Find the area that is common to the polar curves $r = \cos \theta, r = \sin \theta$.

9. Find the area that is inside the polar curve $r = 4 \sin \theta$ and outside the circle $r = 2$.
10. Find the volume that is above the cone $z = \sqrt{x^2 + y^2}$ and below the sphere $x^2 + y^2 + z^2 = 1$.
11. A cylindrical hole of radius a is drilled through a sphere of radius b ($a < b$). Find the volume of the solid that remains.

SOLUTIONS TO ASSIGNMENT #7

1. The iterated integral $I = \int_{x=0}^{x=1} \left(\int_{y=0}^{y=\sqrt{x}} \sin \left(\frac{\pi(y^3 - 3y)}{2} \right) dy \right) dx$ is equal to the double integral $\int \int_R \sin \left(\frac{\pi(y^3 - 3y)}{2} \right) dA$ for a region R in the x, y plane.

- (a) Sketch R .
- (b) Write the integral with the order of integration reversed.
- (c) Compute I .

Solution:

- (a) See diagram at the end.
- (b) $I = \int_{y=0}^{y=1} \left(\int_{x=y^2}^{x=1} \sin \left(\frac{\pi(y^3 - 3y)}{2} \right) dx \right) dy$.
- (c)

$$I = \int_{y=0}^{y=1} \sin \left(\frac{\pi(y^3 - 3y)}{2} \right) (1 - y^2) dy = \frac{2}{3\pi} \cos \left(\frac{\pi(y^3 - 3y)}{2} \right) \Big|_0^1 = -\frac{4}{3\pi}$$

2. Let D be the region bounded by $y = x$ and $y = 6 - x^2$.

- (a) Sketch D .
- (b) Find $\int \int_D x^2 dA$.

Solution:

- (a) See the diagram at the end. Note that $6 - x^2 = x \iff x = -3, 2$.
- (b)

$$\begin{aligned} \int \int_D x^2 dA &= \int_{x=-3}^{x=2} dx \int_{y=x}^{y=6-x^2} x^2 dy = \int_{-3}^2 x^2 (6 - x^2 - x) dx \\ &= -\frac{x^5}{5} \Big|_{-3}^2 - \frac{x^4}{4} \Big|_{-3}^2 + 2x^3 \Big|_{-3}^2 = \frac{125}{4} \end{aligned}$$

3. Let D be the region, described in polar coordinates by, $0 \leq \theta \leq \pi, 0 \leq r \leq 1 + \cos \theta$.

- (a) Sketch D .
- (b) Compute the area of D .
- (c) Find the average value of distances of points in D from the origin.

Solution:

(a) See the diagram at the end.

(b) The area of D is

$$\begin{aligned} A &= \int_{\theta=0}^{\theta=\pi} \int_{r=0}^{r=1+\cos\theta} r dr d\theta = \int_0^\pi \frac{(1+\cos\theta)^2}{2} d\theta \\ &= \frac{1}{2} \int_0^\pi (1+2\cos\theta+\cos^2\theta) d\theta = \frac{1}{2}(\pi + \pi/2) = \frac{3\pi}{4} \end{aligned}$$

(c) By definition, the average value of a function $f(x, y)$ over a domain D is

$$\text{average value} = \frac{1}{\text{area}(D)} \iint_D f(x, y) dx dy.$$

In this case we have

$$\begin{aligned} \text{average value} &= \frac{4}{3\pi} \int_{\theta=0}^{\theta=\pi} \int_{r=0}^{r=1+\cos\theta} r^2 dr = \frac{4}{3\pi} \int_{\theta=0}^{\theta=\pi} \frac{(1+\cos\theta)^3}{3} d\theta \\ &= \frac{4}{9\pi} \int_{\theta=0}^{\theta=\pi} (1+3\cos\theta+3\cos^2\theta+\cos^3\theta) d\theta \\ &= \frac{4}{9\pi}(\pi + 3\pi/2) = \frac{10}{9} \end{aligned}$$

4. Determine the following integrals:

(a) $\iint_D (|x| + |y|) dA$, where D is the region $x^2 + y^2 \leq a^2$ and a is a positive constant.

(b) $\iint_T \sqrt{a^2 - x^2} dA$, where T is the triangle with vertices $(0, 0)$, $(a, 0)$, (a, a) .

(c) $\iint_D \frac{1}{x^2 + y^2} dA$, where D is the region in the first quadrant bounded by

$$y = 0, y = x, x^2 + y^2 = 1/4, x^2 + y^2 = 1.$$

(d) $\iint_R (\sin xy + x^2 - y^2 + 3) dx dy$, where R is the region inside the circle $x^2 + y^2 = a^2$ and outside the circle $x^2 + y^2 = b^2$, and a, b are constants satisfying $0 < b < a$.

Solution:

(a)

$$\begin{aligned} \iint_D (|x| + |y|) dA &= 4 \int_{\theta=0}^{\theta=\pi/2} d\theta \int_{r=0}^{r=a} (r \cos\theta + r \sin\theta) r dr \\ &= \frac{4a^3}{3} \int_{\theta=0}^{\theta=\pi/2} (\cos\theta + \sin\theta) d\theta = \frac{8a^3}{3} \end{aligned}$$

(b)

$$\begin{aligned}\iint_T \sqrt{a^2 - x^2} dA &= \int_{x=0}^{x=a} dx \int_{y=0}^{y=x} \sqrt{a^2 - x^2} dy = \int_{x=0}^{x=a} x\sqrt{a^2 - x^2} dx \\ &= -\frac{1}{3}(a^2 - x^2)^{3/2} \Big|_0^a = \frac{a^3}{3}.\end{aligned}$$

(c) $\iint_D \frac{1}{x^2 + y^2} dA = \int_{\theta=0}^{\theta=\pi/4} \int_{r=1/2}^{r=1} \frac{1}{r} dr d\theta = \frac{\pi \ln 2}{4}.$

(d) By symmetry $\iint_R \sin xy dxdy = 0$ and $\iint_R x^2 dxdy = \iint_R y^2 dxdy$, and therefore $\iint_R (\sin xy + x^2 - y^2 + 3) dxdy = 3 \text{area } R = 3\pi(a^2 - b^2).$

5. Find the volume above the x, y plane, below the surface $z = e^{-(x^2+y^2)}$ and inside the cylinder $x^2 + y^2 = 4$.

Solution: The volume is $V = \int_{\theta=0}^{\theta=2\pi} d\theta \int_{r=0}^{r=2} e^{-r^2} r dr = 2\pi \frac{e^{-r^2}}{-2} \Big|_{r=0}^{r=2} = \pi(1 - e^{-4}).$

6. Find the volume above the x, y plane and below the surface $z = e^{-(x^2+y^2)}$.

Solution: The volume is $V = \int_{\theta=0}^{\theta=2\pi} d\theta \int_{r=0}^{r=\infty} e^{-r^2} r dr = 2\pi \frac{e^{-r^2}}{-2} \Big|_{r=0}^{r=\infty} = \pi.$

7. The iterated integral $\int_{x=0}^{x=4} \left(\int_{y=\sqrt{x}}^{y=2} e^{y^3} dy \right) dx$ can be written in the form $\iint_D e^{y^3} dA$ for a region D .

(a) Sketch D .

(b) Evaluate $\int_{x=0}^{x=4} \left(\int_{y=\sqrt{x}}^{y=2} e^{y^3} dy \right) dx.$

Solution:

(a) See the diagram at the end.

(b) $\int_{x=0}^{x=4} \left(\int_{y=\sqrt{x}}^{y=2} e^{y^3} dy \right) dx = \int_{y=0}^{y=2} dy \int_{x=0}^{x=y^2} e^{y^3} dx = \int_{y=0}^{y=2} y^2 e^{y^3} dy = \frac{e^{y^3}}{3} \Big|_0^2 = \frac{e^8 - 1}{3}$

8. Compute the double integral $\iint_D (x + y) dA$, where D is the domain that lies to the right of the y -axis and between the circles $x^2 + y^2 = 1$, $x^2 + y^2 = 4$.

Solution:

$$\begin{aligned}\iint_D (x + y) dA &= \int_{\theta=-\pi/2}^{\theta=\pi/2} \int_{r=1}^{r=2} (r \cos \theta + r \sin \theta) r dr d\theta \\ &= \frac{7}{3} \int_{\theta=-\pi/2}^{\theta=\pi/2} (\cos \theta + \sin \theta) d\theta = \frac{14}{3}\end{aligned}$$

9. Find the area that is common to the polar curves $r = \cos \theta$, $r = \sin \theta$.

Solution: The area is

$$A = 2 \int_{\theta=0}^{\theta=\pi/4} \int_{r=0}^{r=\sin \theta} r dr d\theta = \int_{\theta=0}^{\theta=\pi/4} \sin^2 \theta d\theta = \pi/8$$

10. Find the area that is inside the polar curve $r = 4 \sin \theta$ and outside the circle $r = 2$.

Solution:

$$\begin{aligned} A &= \int_{\theta=\pi/6}^{\theta=5\pi/6} \int_{r=2}^{r=4 \sin \theta} r dr d\theta = \int_{\theta=\pi/6}^{\theta=5\pi/6} (8 \sin^2 \theta - 2) d\theta \\ &= \int_{\theta=\pi/6}^{\theta=5\pi/6} (4(1 - \cos 2\theta) - 2) d\theta = \int_{\theta=\pi/6}^{\theta=5\pi/6} (2 - 4 \cos 2\theta) d\theta \\ &= \frac{4\pi}{3} - 2 \sin 2\theta \Big|_{\theta=\pi/6}^{\theta=5\pi/6} = \frac{4\pi}{3} - 2 \left(\sin \frac{5\pi}{3} - \sin \frac{2\pi}{3} \right) = \frac{4\pi}{3} + 2\sqrt{3} \end{aligned}$$

11. Find the volume that is above the cone $z = \sqrt{x^2 + y^2}$ and below the sphere $x^2 + y^2 + z^2 = 1$.

Solution:

$$\begin{aligned} V &= \int_{\theta=0}^{\theta=2\pi} \int_{r=0}^{r=1/\sqrt{2}} (\sqrt{1-r^2} - r) r dr d\theta = 2\pi \int_{r=0}^{r=1/\sqrt{2}} (\sqrt{1-r^2} - r) r dr \\ &= 2\pi \left(-\frac{1}{3}(1-r^2)^{3/2} \Big|_{r=0}^{r=1/\sqrt{2}} - \frac{r^3}{3} \Big|_{r=0}^{r=1/\sqrt{2}} \right) = \frac{2\pi}{3}(1 - 1/\sqrt{2}) \end{aligned}$$

12. A cylindrical hole of radius a is drilled through a sphere of radius b ($a < b$). Find the volume of the solid that remains.

Solution:

The volume of the drilled out piece is

$$V = 2 \int_{\theta=0}^{\theta=2\pi} \int_{r=0}^{r=a} \sqrt{b^2 - r^2} r dr d\theta = -\frac{4\pi}{3}(b^2 - r^2)^{3/2} \Big|_{r=0}^{r=a} = \frac{4\pi}{3} (b^3 - (b^2 - a^2)^{3/2})$$

Therefore the volume of the remaining piece is $\frac{4\pi}{3} (b^2 - a^2)^{3/2}$.




Figure 1: Question 1(a), $0 \leq x \leq 1, 0 \leq y \leq \sqrt{x}$




Figure 2: Question 2(a), the region bounded by $y = x, y = 6 - x^2$




Figure 3: Question 3(a), $0 \leq \theta \leq \pi, 0 \leq r \leq 1 + \cos \theta$




Figure 4: Question 7(a), $0 \leq x \leq 4, \sqrt{x} \leq y \leq 2$